


Recent results on anomalous J/ψ suppression in Pb-Pb collisions at 158 AGeV/c

Helena Santos for the NA50 Collaboration

B. Alessandro¹¹, C. Alexa⁴, R. Arnaldi¹¹, M. Atayan¹³, C. Baglin², A. Baldit³, M. Bedjidian¹², S. Beolè¹¹, V. Boldea⁴, P. Bordalo^{7,a}, S.R. Borenstein^{10,b}, G. Borges⁷, A. Bussière², L. Capelli¹², C. Castanier³, J. Castor³, B. Chaurand¹⁰, B. Cheynis¹², E. Chiavassa¹¹, C. Cicalò⁵, T. Claudino⁷, M.P. Comets⁹, S. Constantinescu⁴, P. Cortese¹, J. Cruz⁷, A. DeFalco⁵, N. DeMarco¹¹, G. Dellacasa¹, A. Devaux³, S. Dita⁴, O. Drapier¹⁰, B. Espagnon³, J. Fargeix³, P. Force³, M. Gallio¹¹, Y.K. Gavrilo⁸, C. Gerschel⁹, P. Giubellino^{11,c}, M.B. Golubeva⁸, M. Gonin¹⁰, A.A. Grigorian¹³, S. Grigorian¹³, J.Y. Grossiord¹², F.F. Guber⁸, A. Guichard¹², H. Gulkanyan¹³, R. Hakobyan¹³, R. Haroutunian¹², M. Idzik^{11,d}, D. Jouan⁹, T.L. Karavitcheva⁸, L. Kluberg¹⁰, A.B. Kurepin⁸, Y. Le Bornec⁹, C. Lourenço⁶, P. Macciotta⁵, M. Mac Cormick⁹, A. Marzari-Chiesa¹¹, M. Masera¹¹, A. Masoni⁵, M. Monteno¹¹, A. Musso¹¹, P. Petiau¹⁰, A. Piccotti¹¹, J.R. Pizzi¹², W.L. Prado da Silva^{11,e}, F. Prino¹, G. Puddu⁵, C. Quintans⁷, L. Ramello¹, S. Ramos^{7,a}, P. Rato Mendes⁷, L. Riccati¹¹, A. Romana¹⁰, H. Santos⁷, P. Saturnini³, E. Scalas¹, E. Scomparin¹¹, S. Serchi⁵, R. Shahoyan^{7,f}, F. Sigaud¹¹, M. Sitta¹, P. Sonderegger^{6,a}, X. Tarrago⁹, N.S. Topilskaya⁸, G.L. Usai⁵, E. Vercellin¹¹, L. Villatte⁹, N. Willis⁹ and T. Wu⁹

¹ Università del Piemonte Orientale, Alessandria and INFN-Torino, Italy ² LAPP, CNRS-IN2P3, Annecy-le-Vieux, France ³ LPC, Univ. Blaise Pascal and CNRS-IN2P3, Aubière, France ⁴ IFA, Bucharest, Romania ⁵ Università di Cagliari/INFN, Cagliari, Italy ⁶ CERN, Geneva, Switzerland ⁷ LIP, Lisbon, Portugal ⁸ INR, Moscow, Russia ⁹ IPN, Univ. de Paris-Sud and CNRS-IN2P3, Orsay, France ¹⁰ Laboratoire Leprince-Ringuet, Ecole Polytechnique and CNRS-IN2P3, Palaiseau, France ¹¹ Università di Torino/INFN, Torino, Italy ¹² IPN, Univ. Claude Bernard Lyon-I and CNRS-IN2P3, Villeurbanne, France

a) also at IST, Universidade Técnica de Lisboa, Lisbon, Portugal, b) on leave of absence from York College, CUNY, New York, USA, c) also at CERN, Geneva, Switzerland, d) also at Faculty of Physics and Nuclear Techniques, University of Mining and Metallurgy, Cracow, Poland, e) now at UERJ, Rio de Janeiro, Brazil, f) on leave of absence from YerPhI, Yerevan, Armenia



We report our latest results on charmonium suppression as measured at the CERN-SPS in Pb-Pb interactions at an incident beam momentum of 158 GeV/c per nucleon. Preliminary results obtained from the most recent sample of data collected in year 2000 under improved experimental conditions are compared with published results. For the most peripheral Pb-Pb collisions, J/ψ suppression agrees with the normal absorption measured from interactions of lighter nuclei while a steady increasing abnormal suppression is observed with increasing centrality.

1 Introduction

The study of charmonium production in ultrarelativistic heavy ion collisions by experiments NA38 and NA50 has been motivated by the search for a phase transition of nuclear matter from its normal state to a deconfined quark gluon plasma, as predicted to occur, under extreme energy densities or temperature conditions, by non-perturbative QCD. In such a deconfined matter, the J/ψ is expected to be suppressed by Debye colour screening¹. As the vector-meson is produced at the very early stages of the collision, its leptonic decay provides a probe of these stages, since this mode is insensitive to strong interactions and, therefore, to the hadronization stage.

2 Data selection and analysis method

The main element of the NA50 detector is a muon spectrometer. It is complemented by centrality detectors which measure the neutral transverse energy E_T , the beam spectators energy E_{ZDC} and the charged multiplicity produced in the interaction. Several beam monitoring devices are also used for various purposes. A detailed description can be found in². The goal of the year 2000 Pb-Pb data collection is to investigate, in particular, the most peripheral collisions which, in previous data samples, could be contaminated by Pb-air interactions. The data have been taken, therefore, with the target placed in the vacuum pipe.

The event selection is done according to the following criteria: dimuons are selected in the rapidity window $2.92 \leq y_{lab} < 3.92$ ($0 \leq y_{CM} < 1$) and with a Collins-Soper angle $|\cos\theta_{CS}| < 0.5$, leading to an acceptance of around 14% in the mass region of interest. Parasitic interactions of an incident Pb-ion in the Beam Hodoscope and upstream from the target are rejected by a BH interaction detector and by Anti-halo counters. On-target interactions are selected requiring a good correlation between hits in the two planes of the Multiplicity Detector. Double interactions are discarded by a shape analysis of the signal in the Electromagnetic Calorimeter. Residual pileup events are rejected by a 2σ cut in the $E_T - E_{ZDC}$ correlation³. Off-target muons are excluded by a cut on the transverse distance between the extrapolated muon track and the beam line.

In each centrality bin of the analysis, the opposite-sign dimuon mass spectrum is adjusted in the pertinent mass region, accounting for the different ingredients: J/ψ , ψ' and the continuum formed by Drell-Yan, $D\bar{D}$ and the combinatorial background from π and K decays. The shapes of the resonances are obtained by Monte Carlo simulation followed by reconstruction with the same program as used for the real data analysis. Drell-Yan and $D\bar{D}$ are generated at leading order by Pythia and treated by the NA50 simulation-reconstruction chain. The combinatorial background is properly extracted from like-sign pairs through the formula $N_{BG} = 2\sqrt{N^{++}N^{--}}$.

3 Before the year 2000 – The state of the art

The NA38/NA51/NA50 experiments have verified that, as expected theoretically, the Drell-Yan cross-section $\sigma(DY)$ is proportional to the number of elementary nucleon-nucleon collisions, from proton-proton up to lead-lead interactions included (see Fig. 1). The ratio of the cross-sections for J/ψ and Drell-Yan production is therefore proportional to the J/ψ production cross-section per nucleon-nucleon collision. Moreover, the ratio $B_{\mu\mu}\sigma(J/\psi)/\sigma(DY)$ is directly measured in the so-called standard analysis method and insensitive to reconstruction or trigger inefficiencies (identical muon pair topology for J/ψ and Drell-Yan events) or to absolute normalization uncertainties which, obviously, cancel out. Fig. 2 shows the ratio $B_{\mu\mu}\sigma(J/\psi)/\sigma(DY)$ as a function of E_T , for the previously collected Pb-Pb data samples – 1995^{2,4}, 1996⁵ and 1998⁶. The closed points are the results obtained with the standard analysis method described above while the open points are the 1996 and 1998 results obtained through the "Minimum Bias analysis" method explained in [4,5]. The continuous line is the J/ψ normal absorption in nuclear

matter and fits the NA51 p-p and p-d and NA38 p-A and S-U results. A clear departure of the $B_{\mu\mu}\sigma(J/\psi)/\sigma(DY)$ ratios from this normal absorption curve is observed at mid-centrality, suggesting the onset of an anomalous J/ψ suppression mechanism. Furthermore, no saturation of the ratio at high E_T is observed, as opposed to the absorption curve trend.




Figure 1: Drell-Yan K-factor as a function of $A_{\text{projectile}} \times B_{\text{target}}$




Figure 2: $J/\psi/DY$ as a function of E_T ; the absorption curve is a fit to the NA38 p-A and S-U data

4 The preliminary year 2000 results

The NA50 Collaboration has a new high statistics p-A data sample, obtained with a 450 GeV/c incident energy beam. Since the J/ψ absorption cross sections for p-A and S-U collisions are compatible within the statistical errors, a simultaneous Glauber fit is performed which leads to the common value of $\sigma(J/\psi)_{\text{abs}} = 4.3 \pm 0.5 \text{ mb}$ ⁷. This result, combined with the NA38 S-U data reanalyzed in order to have the same analysis conditions, leads to an updated normal nuclear absorption.

From the above described fit to the dimuon invariant mass spectra, we extract the J/ψ and the Drell-Yan yields (the latter in the mass interval 2.9 - 4.5 GeV/c²) and obtain the ratio $B_{\mu\mu}\sigma(J/\psi)/\sigma(DY)$ and its dependence as a function of E_T or E_{ZDC} . Fig. 3 shows the agreement among independent analyses with slightly different data selections and fit methods. It also shows that peripheral collisions indeed follow the absorption curve. On the other hand, the Pb-Pb 2000 data confirm the departure from this normal behaviour of the $J/\psi/DY$ ratios at mid-centrality, already observed in the previously collected data. The results also confirm the steady decrease of the J/ψ production rate at high E_T . The same J/ψ suppression pattern is seen in Fig. 4 when E_{ZDC} is used as the centrality estimator. For this variable, the agreement is quite good between new and already published results, except for the most peripheral collisions, probably due to Pb-air interactions in the 1996 setup conditions.

In the 2.9-4.5 GeV/c² invariant mass range, muons from J/ψ and Drell-Yan have similar features, namely acceptance and momentum. Nevertheless, different PDFs chosen to generate Drell-Yan lead to different mass distributions, as shown in Fig. 5. The difference under the J/ψ peak between the Drell-Yan cross sections generated with MRS43 and GRV LO amounts to 10% systematic effect in the $J/\psi/DY$ normalizations, independent of centrality. To overcome this uncertainty, the 4.2-7.0 GeV/c² mass range is chosen as the new J/ψ suppression reference. In fact, beyond 4.2 GeV/c², the Drell-Yan pairs are the only contribution to the dimuon mass spectrum and thus both structure functions give the same normalization after adjustment to the data.

Fig. 6 shows all NA50 results using the new mass range for Drell-Yan as reference. All data sets agree, regardless on the fact that Pb-Pb 1995 results were obtained using GRV LO PDFs while 1996, 1998 and 2000 used MRS43. Pb-Pb 1998 and 2000 analyses with GRV LO is also shown.




Figure 3: $J/\psi/DY$ as a function of E_T




Figure 4: $J/\psi/DY$ as a function of E_{ZDC}




Figure 5: The Drell-Yan mass distributions for MRS43 and GRV LO PDFs




Figure 6: $J/\psi/DY$ as a function of E_T

5 Conclusions

In the most recently collected Pb-Pb data sample, the NA50 Collaboration has measured $B_{\mu\mu}\sigma(J/\psi)/\sigma(DY)$ as a function of centrality. With respect to previously collected data samples, the use of a setup with the target region under vacuum allows to extend, confidently, the analysis to very peripheral collisions.

These new data, confirming previous analyses results, show that J/ψ production follows the normal nuclear absorption pattern for peripheral Pb-Pb collisions, while exhibiting a clear departure for central collisions and suggesting, therefore, the onset of another J/ψ suppression mechanism.

Acknowledgments

This work was partially supported by Fundação para a Ciência e a Tecnologia.

References

1. T. Matsui and H. Satz, Phys. Lett. B178 (1986) 416.
2. M.C. Abreu et al. (NA50 Coll.), Phys. Lett. B410 (1997) 327.
3. C. Quintans, PhD Thesis, Instituto Superior Técnico, Lisboa (2002).
4. M.C. Abreu et al. (NA50 Coll.), Phys. Lett. B410 (1997) 337.
5. M.C. Abreu et al. (NA50 Coll.), Phys. Lett. B450 (1999) 456.
6. M.C. Abreu et al. (NA50 Coll.), Phys. Lett. B477 (2000) 28.

7. B. Alessandro et al. (NA50 Coll.), Phys. Lett. B553 (2003) 167.